

Example Sheet 2

1. A particle of mass m is confined to a one-dimensional box $0 \leq x \leq a$ (the potential $V(x)$ is zero inside the box, and infinite outside). Show that the energy eigenvalues are $E_n = \hbar^2 \pi^2 n^2 / 2ma^2$ for $n = 1, 2, \dots$, and determine corresponding normalised energy eigenstates $\psi_n(x)$. Show that the expectation value and the uncertainty for a measurement of \hat{x} in the state ψ_n are given by

$$\langle \hat{x} \rangle_n = \frac{a}{2} \quad \text{and} \quad (\Delta x)_n^2 = \frac{a^2}{12} \left(1 - \frac{6}{\pi^2 n^2} \right).$$

Does the limit $n \rightarrow \infty$ agree with what you would expect for a classical particle in this potential?

2. Write down the Hamiltonian H for a harmonic oscillator of mass m and frequency ω . Express $\langle H \rangle$ in terms of $\langle \hat{x} \rangle$, $\langle \hat{p} \rangle$, Δx and Δp , all defined for some normalised state ψ . Use the Uncertainty Relation to deduce that $E \geq \frac{1}{2} \hbar \omega$ for any energy eigenvalue E .

3. Let $\Psi(x, t)$ be a solution of the time-dependent Schrödinger Equation with zero potential (corresponding to a free particle). Show that

$$\Phi(x, t) = \Psi(x - ut, t) e^{ikx} e^{-i\omega t}$$

is also a solution if the constants k and ω are chosen suitably, in terms of u . Express $\langle \hat{x} \rangle_\Phi$ and $\langle \hat{p} \rangle_\Phi$ in terms of $\langle \hat{x} \rangle_\Psi$ and $\langle \hat{p} \rangle_\Psi$. Are the results consistent with Ehrenfest's Theorem?

4. The energy levels of the harmonic oscillator are $E_n = (n + \frac{1}{2}) \hbar \omega$ for $n = 0, 1, 2, \dots$ and the corresponding stationary state wavefunctions are

$$\psi_n(x) = h_n(y) e^{-y^2/2} \quad \text{where} \quad y = (m\omega/\hbar)^{1/2} x$$

and h_n is a polynomial of degree n with $h_n(-y) = (-1)^n h_n(y)$. Using *only* the orthogonality relations

$$(\psi_m, \psi_n) = \delta_{mn},$$

determine ψ_2 and ψ_3 up to an overall constant in each case.

Give an expression for the quantum state of the oscillator $\Psi(x, t)$ if the initial state is $\Psi(x, 0) = \sum_{n=0}^{\infty} c_n \psi_n(x)$, where c_n are complex constants. Deduce that

$$|\Psi(x, 2p\pi/\omega)|^2 = |\Psi(-x, (2q+1)\pi/\omega)|^2$$

for any integers $p, q \geq 0$. Comment on this result, considering the particular case in which $\Psi(x, 0)$ is sharply peaked around position $x = a$.

5. Consider the Schrödinger Equation in one dimension with potential $V(x)$. Show that for a stationary state, the probability current J is independent of x .

Now suppose that an energy eigenstate $\psi(x)$ corresponds to scattering by the potential and that $V(x) \rightarrow 0$ as $x \rightarrow \pm\infty$. Given the asymptotic behaviour

$$\psi(x) \sim e^{ikx} + B e^{-ikx} \quad (x \rightarrow -\infty) \quad \text{and} \quad \psi(x) \sim C e^{ikx} \quad (x \rightarrow +\infty)$$

show that $|B|^2 + |C|^2 = 1$. How should this be interpreted?

6. A particle is incident on a potential barrier of width a and height U . Assuming that $U = 2E$, where $E = \hbar^2 k^2 / 2m$ is the kinetic energy of the incident particle, find the transmission probability. [*Work through the algebra, which simplifies in this case, rather than quoting the general result.*]

7. Consider the time-independent Schrödinger Equation with potential $V(x) = -U\delta(x)$. Show that there is a scattering solution with energy eigenvalue $E = \hbar^2 k^2/2m$ for any real $k > 0$ and find the transmission and reflection amplitudes $A_{\text{tr}}(k)$ and $A_{\text{ref}}(k)$. [Recall from Example 9 on Sheet 1 that the wavefunction ψ is continuous, but satisfies $\psi'(0+) - \psi'(0-) = -(2mU/\hbar^2)\psi(0)$.]

Is the solution above still an eigenfunction of the Hamiltonian if k is allowed to take complex values? Show that $A_{\text{tr}}(k)$ and $A_{\text{ref}}(k)$ are singular at $k = i\kappa$ for a certain real, positive value of κ . By first re-scaling the scattering solution, find a bound state (normalisable) solution in the potential. What is the energy of this bound state?

8. A particle of mass m is in a one-dimensional infinite square well (a potential box) with $V = 0$ for $0 < x < a$ and $V = \infty$ otherwise. The normalised wavefunction for the particle at time $t = 0$ is

$$\Psi(x, 0) = Cx(a - x) .$$

(i) Determine the real constant C .

(ii) By expanding $\Psi(x, 0)$ as a linear combination of energy eigenfunctions (found in Example 1 above), obtain an expression for $\Psi(x, t)$, the wavefunction at time t .

(iii) A measurement of the energy is made at time $t > 0$. Show that the probability that this yields the result $E_n = \hbar^2 \pi^2 n^2 / 2ma^2$ is $960/\pi^6 n^6$ if n is odd, and zero if n is even. Why should the result for n even be expected? Which value of the energy is most likely, and why is its probability so close to unity?

9. A quantum system has Hamiltonian H with normalised eigenstates ψ_n and corresponding energies E_n ($n = 1, 2, 3, \dots$). A linear operator Q is defined by its action on these states:

$$Q\psi_1 = \psi_2 , \quad Q\psi_2 = \psi_1 , \quad Q\psi_n = 0 \quad n > 2 .$$

Show that Q has eigenvalues ± 1 (in addition to zero) and find the corresponding normalised eigenstates χ_{\pm} , in terms of energy eigenstates. Calculate $\langle H \rangle$ in each of the states χ_{\pm} .

A measurement of Q is made at time zero, and the result $+1$ is obtained. The system is then left undisturbed for a time t , at which instant another measurement of Q is made. What is the probability that the result will again be $+1$? Show that the probability is zero if the measurement is made when a time $T = \pi\hbar/(E_2 - E_1)$ has elapsed (assume $E_2 - E_1 > 0$).

10. In the previous example, suppose that an experimenter makes n successive measurements of Q at regular time intervals T/n . If the result $+1$ is obtained for one measurement, show that the amplitude for the next measurement to give $+1$ is

$$A_n = 1 - \frac{iT(E_1 + E_2)}{2\hbar n} + \mathcal{O}\left(\frac{1}{n^2}\right) .$$

The probability that all n measurements give the result $+1$ is then $P_n = (|A_n|^2)^n$. Show that

$$\lim_{n \rightarrow \infty} P_n = 1 .$$

Interpreting χ_{\pm} as the 'not-boiling' and 'boiling' states of a two-state 'quantum pot', this shows that a watched quantum pot never boils (also called the Quantum Zeno Paradox).

11. Let H be a Hamiltonian and ψ any normalised eigenstate with energy E . Show that, for any operator A ,

$$\langle [H, A] \rangle_{\psi} = 0 .$$

For a particle in one dimension, let $H = T + V$ where $T = \hat{p}^2/2m$ is the kinetic energy and $V(\hat{x})$ is any (real) potential. By setting $A = \hat{x}$ in the result above and using the canonical commutation relation between position and momentum, show that $\langle \hat{p} \rangle_{\psi} = 0$.

Now assume further that $V(\hat{x}) = k\hat{x}^n$ (with k and n constants). By taking $A = \hat{x}\hat{p}$, show that

$$\langle T \rangle_{\psi} = \frac{n}{n+2} E \quad \text{and} \quad \langle V \rangle_{\psi} = \frac{2}{n+2} E .$$